

# Hardy-Littlewood Conjecture and Exceptional real Zero

JinHua Fei

ChangLing Company of Electronic Technology Baoji Shannxi P.R.China

E-mail: feijinhuaoujian@msn.com

**Abstract.** In this paper, we assume that Hardy-Littlewood Conjecture, we got a better upper bound of the exceptional real zero for a class of module.

**Keyword.** Hardy-Littlewood Conjecture, Exceptional real Zero

**MR(2000) Subject Classification** 11P32, 11M20

In this paper, we reveal the relationship between Hardy-Littlewood Conjecture and the exceptional real zero. Assume that Hardy-Littlewood Conjecture, we obtained a good results of the exceptional real zero.

In this paper, I generalize the results of my paper "An Application of Hardy-Littlewood Conjecture".  $q$  is the prime number and  $q \equiv 3 \pmod{4}$  is improved to  $q$  is odd square-free and  $q \equiv 3 \pmod{4}$ . we know that the module  $q$  of the exceptional primitive real character are square-free when  $q$  is odd positive integer.

First, we give Hardy-Littlewood Conjecture.

**Hardy-Littlewood Conjecture.** When  $N$  is even integer and  $N \geq 6$ , we have

$$\sum_{\substack{3 \leq p_1, p_2 \leq N \\ p_1 + p_2 = N}} 1 \approx \frac{N}{\varphi(N)} \prod_{p \nmid N} \left(1 - \frac{1}{(p-1)^2}\right) \frac{N}{\log^2 N}$$

where  $p_1, p_2, p$  are the prime numbers,  $\varphi(n)$  is Euler function.

Under the above conjecture, we have the following theorem

**Theorem.** Let  $q$  is odd square-free and  $q \equiv 3 \pmod{4}$ , it has exceptional real character  $\chi$ , and its Dirichlet  $L(s, \chi)$  function has an exceptional real zero  $\beta$ . If Hardy-Littlewood Conjecture is correct, then there is a positive constant  $c$ , we have

$$\beta \leq 1 - \frac{c}{\log^2 q}$$

Now, we do some preparation work.

**Lemma 1 .** Assuming that the Hardy-Littlewood Conjecture. Let  $n$  is any positive integer,  $q$  is odd integer and large sufficiently, then

$$\sum_{\substack{3 \leq p_1, p_2 \leq 2nq \\ p_1 + p_2 = 2nq}} 1 \geq \frac{q}{\varphi(q)} \frac{2nqd}{(\log 2nq)^2}$$

where

$$d = \prod_{3 \leq p} \left( 1 - \frac{1}{(p-1)^2} \right)$$

and  $p_1, p_2, p$  are the prime numbers.

**Proof .** By Hardy-Littlewood Conjecture, when  $N$  is large sufficiently, we have

$$\sum_{\substack{3 \leq p_1, p_2 \leq N \\ p_1 + p_2 = N}} 1 \geq \frac{N}{2\varphi(N)} \prod_{p \nmid N} \left( 1 - \frac{1}{(p-1)^2} \right) \frac{N}{\log^2 N}$$

because

$$\prod_{p \nmid N} \left(1 - \frac{1}{(p-1)^2}\right) \geq \prod_{3 \leq p} \left(1 - \frac{1}{(p-1)^2}\right)$$

and

$$\frac{2nq}{\varphi(2nq)} = \prod_{p|2nq} \frac{p}{p-1} \geq \prod_{p|2q} \frac{p}{p-1} = \frac{2q}{\varphi(q)}$$

We choose  $N = 2nq$  , This completes the proof of Lemma 1 .

**Lemma 2.** Let  $m$  is positive integer and  $n$  is integer, then

$$\sum_{k=1}^m e\left(\frac{kn}{m}\right) = \begin{cases} m & \text{if } n \equiv 0 \pmod{m} \\ 0 & \text{otherwise} \end{cases}$$

where  $e(x) = e^{2\pi i x}$ . The lemma 2 is obvious

**Lemma 3.** Let  $c_1$  be the positive constant. if  $(a, q) = 1$ , then

$$\pi(x; q, a) = \frac{Lix}{\varphi(q)} - \frac{\chi(a)}{\varphi(q)} \int_2^x \frac{u^{\beta-1}}{\log u} du + O\left(x \exp(-c_1 \sqrt{\log x})\right)$$

when there is an exceptional character  $\chi$  modulo  $q$  and  $\beta$  is the concomitant zero. Where  $Lix = \int_2^x \frac{du}{\log u}$  and  $\exp(x) = e^x$

The lemma 3 follows from the References [2], Corollary 11.20 of the page 381

It is easy to see that

$$Lix = \int_2^x \frac{du}{\log u} = \frac{x}{\log x} + O\left(\frac{x}{\log^2 x}\right)$$

and

$$\int_2^x \frac{u^{\beta-1}}{\log u} du = \frac{x^\beta}{\beta \log x} + O\left(\frac{x^\beta}{\log^2 x}\right)$$

**Lemma 4.** if  $\chi$  is a primitive character modulo  $m$ , then

$$\sum_{k=1}^m \chi(k) e\left(\frac{nk}{m}\right) = \bar{\chi}(n) \tau(\chi)$$

where  $\tau(\chi) = \sum_{k=1}^m \chi(k) e\left(\frac{k}{m}\right)$ .

The lemma 4 follows from the References [1], the page 47.

**Lemma 5.** if  $m$  is odd square-free and  $\chi$  is a primitive real character modulo  $m$ , then

$$\tau(\chi) = \begin{cases} \sqrt{m} & \text{if } m \equiv 1 \pmod{4} \\ i\sqrt{m} & \text{if } m \equiv 3 \pmod{4} \end{cases}$$

The lemma 5 follows from the References [1], the theorem 3.3 of the page 49.

**Lemma 6.** We give the value of two sums, they are used in the proof of the Theorem.

(1)

$$\sum_{k=1}^q \left( \sum_{\substack{a=1 \\ (a,q)=1}}^{q-1} e\left(\frac{ak}{q}\right) \right)^2 = \sum_{k=1}^q \sum_{\substack{a=1 \\ (a,q)=1}}^{q-1} e\left(\frac{ak}{q}\right) \sum_{\substack{b=1 \\ (b,q)=1}}^{q-1} e\left(\frac{bk}{q}\right)$$

$$= \sum_{k=1}^q \sum_{\substack{a=1 \\ (a,q)=1}}^{q-1} \sum_{\substack{b=1 \\ (b,q)=1}}^{q-1} e\left(\frac{(a+b)k}{q}\right) = \sum_{\substack{a=1 \\ (a,q)=1}}^{q-1} \sum_{\substack{b=1 \\ (b,q)=1}}^{q-1} \sum_{k=1}^q e\left(\frac{(a+b)k}{q}\right)$$

$$= q \sum_{\substack{a=1 \\ (a,q)=1}}^{q-1} \sum_{\substack{b=1 \\ (b,q)=1, a+b=q}}^{q-1} 1 = q \varphi(q)$$

(2)

$$\sum_{k=1}^q \chi(k) \sum_{\substack{a=1 \\ (a,q)=1}}^{q-1} e\left(\frac{ak}{q}\right) = \sum_{\substack{a=1 \\ (a,q)=1}}^{q-1} \sum_{k=1}^q \chi(k) e\left(\frac{ak}{q}\right) = \tau(\chi) \sum_{\substack{a=1 \\ (a,q)=1}}^{q-1} \bar{\chi}(a) = 0$$

where  $\chi$  is the primitive character modulo  $q$ ,

This completes the proof of Lemma 6 .

#### **PROOF OF THEOREM.**

*The first part.*

By Lemma 2, when  $x \geq q^4$ , we have

$$\sum_{k=1}^q \left( \sum_{3 \leq p \leq x} e\left(\frac{kp}{q}\right) \right)^2 = \sum_{k=1}^q \sum_{3 \leq p_1 \leq x} \sum_{3 \leq p_2 \leq x} e\left(\frac{k(p_1 + p_2)}{q}\right)$$

$$= \sum_{3 \leq p_1 \leq x} \sum_{3 \leq p_2 \leq x} \sum_{k=1}^q e\left(\frac{k(p_1 + p_2)}{q}\right) = q \sum_{\substack{3 \leq p_1, p_2 \leq x \\ p_1 + p_2 \equiv 0 \pmod{q}}} 1 \geq q \sum_{n=1}^{\lfloor \frac{x}{2q} \rfloor} \sum_{\substack{3 \leq p_1, p_2 \leq x \\ p_1 + p_2 = 2nq}} 1$$

by Lemma 1 , the above formula

$$\begin{aligned} &\geq \frac{q^2}{\varphi(q)} \sum_{n=1}^{\lfloor \frac{x}{2q} \rfloor} \frac{2nqd}{\log^2 2nq} \geq \frac{q^2}{\varphi(q)} \sum_{n=1}^{\lfloor \frac{x}{2q} \rfloor} \frac{2nqd}{\log^2 x} \geq \frac{2dq^3}{\varphi(q) \log^2 x} \sum_{n=1}^{\lfloor \frac{x}{2q} \rfloor} n \\ &= \frac{2dq^3}{\varphi(q) \log^2 x} \cdot \frac{\lfloor \frac{x}{2q} \rfloor (\lfloor \frac{x}{2q} \rfloor + 1)}{2} \geq \frac{q d x^2}{4\varphi(q) \log^2 x} + O\left(\frac{xq^2}{\varphi(q) \log^2 x}\right) \end{aligned}$$

*The second part.*

When  $1 \leq k \leq q$ , we have

$$\sum_{3 \leq p \leq x} e\left(\frac{pk}{q}\right) = \sum_{\substack{3 \leq p \leq x \\ (p,q)=1}} e\left(\frac{pk}{q}\right) + O(\log q) = \sum_{\substack{a=1 \\ (a,q)=1}}^q e\left(\frac{ak}{q}\right) \sum_{\substack{3 \leq p \leq x \\ p \equiv a \pmod{q}}} 1 + O(\log q)$$

by Lemma 3 and Lemma 4, the above formula

$$= \sum_{\substack{a=1 \\ (a,q)=1}}^q e\left(\frac{ak}{q}\right) \left( \frac{Lix}{\varphi(q)} - \frac{\chi(a)}{\varphi(q)} \int_2^x \frac{u^{\beta-1}}{\log u} du + O\left(x \exp(-c_1 \sqrt{\log x})\right) \right) + O(\log q)$$

$$= \frac{Lix}{\varphi(q)} \sum_{\substack{a=1 \\ (a,q)=1}}^q e\left(\frac{ak}{q}\right) - \frac{\tau(\chi)\chi(k)}{\varphi(q)} \int_2^x \frac{u^{\beta-1}}{\log u} du + O\left(qx \exp(-c_1 \sqrt{\log x})\right)$$

where  $\chi$  is the exceptional primitive real character modulo  $q$ .

therefore

$$\left(\sum_{3 \leq p \leq x} e\left(\frac{pk}{q}\right)\right)^2 = \left(\frac{Lix}{\varphi(q)} \sum_{\substack{a=1 \\ (a,q)=1}}^q e\left(\frac{ak}{q}\right)\right)^2$$

$$- \frac{2\tau(\chi)\chi(k)Lix}{\varphi^2(q)} \left(\sum_{\substack{a=1 \\ (a,q)=1}}^q e\left(\frac{ak}{q}\right)\right) \int_2^x \frac{u^{\beta-1}}{\log u} du + \left(\frac{\chi(k)\tau(\chi)}{\varphi(q)} \int_2^x \frac{u^{\beta-1}}{\log u} du\right)^2$$

$$+ O\left(q^2 x^2 \exp(-c_2 \sqrt{\log x})\right)$$

By Lemma 5 and Lemma 6, we have

$$\sum_{k=1}^q \left(\sum_{3 \leq p \leq x} e\left(\frac{pk}{q}\right)\right)^2 = \frac{q}{\varphi(q)} (Lix)^2 - \frac{q}{\varphi(q)} \left(\int_2^x \frac{u^{\beta-1}}{\log u} du\right)^2$$

$$+O\left(q^3 x^2 \exp(-c_2 \sqrt{\log x})\right)$$

$$= \frac{q x^2}{\varphi(q) \log^2 x} - \frac{q x^{2\beta}}{\varphi(q) \beta^2 \log^2 x} + O\left(\frac{q x^2}{\varphi(q) \log^3 x} + q^3 x^2 \exp(-c_2 \sqrt{\log x})\right)$$

We synthesize the first part and second part, we have

$$\frac{q d x^2}{4 \varphi(q) \log^2 x} \leq \frac{q x^2}{\varphi(q) \log^2 x} - \frac{q x^{2\beta}}{\varphi(q) \beta^2 \log^2 x}$$

$$+O\left(\frac{x^2}{\log^3 x} + \frac{x q^2}{\varphi(q) \log^2 x} + q^3 x^2 \exp(-c_2 \sqrt{\log x})\right)$$

$$\frac{d x^2}{4 \log^2 x} \leq \frac{x^2}{\log^2 x} - \frac{x^{2\beta}}{\beta^2 \log^2 x} + O\left(\frac{x^2}{\log^3 x} + \frac{x q}{\log^2 x} + q^3 x^2 \exp(-c_2 \sqrt{\log x})\right)$$

$$\frac{d}{4} \leq 1 - \frac{x^{2\beta-2}}{\beta^2} + O\left(\frac{1}{\log x} + \frac{q}{x} + q^3 \exp(-c_3 \sqrt{\log x})\right)$$

we take  $\log x = (\frac{4}{c_3} \log q)^2$ , then

$$x^{2\beta-2} \leq 1 - \frac{d}{4} + \frac{c_4}{\log^2 q}$$

we take  $\log q \geq \sqrt{\frac{8c_4}{d}}$ , then

$$x^{2\beta-2} \leq 1 - \frac{d}{8}$$

$$\beta - 1 \leq \frac{\log(1 - \frac{d}{8})}{2 \log x} = -\frac{\log(\frac{8}{8-d})}{2 \log x}$$

therefore

$$\beta \leq 1 - \frac{c}{\log^2 q}$$

This completes the proof of Theorem.

Because  $\tau(\chi_4) = 2i$  and  $\tau(\chi_8) = 2\sqrt{2}$ , when  $q$  is odd square-free, by same method, for module  $4q, q \equiv 1(mod 4)$  and module  $8q, q \equiv 3(mod 4)$ , we have the same conclusion.

## REFERENCES

- [1] Henryk Iwaniec, Emmanuel Kowalski, *Analytic Number Theory*, American mathematical Society, 2004.
- [2] Hugh L. Montgomery, Robert C. Vaughan, *Multiplicative Number Theory I. Classical Theory*, Cambridge University Press, 2006.